




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 498/04, 405/14, 405/12 A61K 31/535, 31/445 // C07D 498/04, 265:00, 209:00	A1	(11) International Publication Number: WO 94/10174 (43) International Publication Date: 11 May 1994 (11.05.94)
(21) International Application Number: PCT/EP93/03054 (22) International Filing Date: 2 November 1993 (02.11.93) (30) Priority data: 9223155.4 5 November 1992 (05.11.92) GB 9309644.4 11 May 1993 (11.05.93) GB 9315202.3 22 July 1993 (22.07.93) GB (71) Applicant (for all designated States except US): SMITH-KLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9E EP (GB). (72) Inventors; and (75) Inventors/Applicants (for US only) : GASTER, Laramie, Mary [GB/GB]; WYMAN, Paul, Adrian [GB/GB]; SmithKline Beecham Pharmaceuticals, Coldharbour Road, The Pinnacles, Harlow, Essex CM19 5AD (GB).		(74) Agents: JONES, Pauline et al.; SmithKline Beecham, Corporate Intellectual Property, Great Burgh, Yew Tree Bottom Road, Epsom, Surrey KT18 5XQ (GB). (81) Designated States: AU, CA, JP, KR, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>

(54) Title: PIPERIDINE DERIVATIVES AS 5-HT₄ RECEPTOR ANTAGONISTS

**(57) Abstract**

Compounds of formula (I) and pharmaceutically acceptable salts thereof, and the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof: X-CO-Y-Z, wherein X is a monocyclic or polycyclic aromatic group, Y is O or NH; Z is of sub-formula (II), wherein -(CH₂)_{n¹} is attached at carbon; and n¹ is 0, 1, 2, 3 or 4; q is 0, 1, 2 or 3; R_a is straight or branched chain alkylene of chain length 1-6 carbon atoms terminally substituted by R₇ wherein R₇ is aryl, 3 to 8 membered cycloalkyl, 3 to 8 membered heterocyclyl, 5 or 6 membered monocyclic heteroaryl or 9 or 10 membered fused bicyclic heteroaryl linked through carbon, or R₇ is C₂₋₇ alkoxy carbonyl or secondary or tertiary hydroxy substituted C₁₋₆ alkyl; and R₆ is hydrogen or C₁₋₆ alkyl; or a compound of formula (I) wherein the CO-Y linkage is replaced by a heterocyclic bioisostere; and their use as pharmaceuticals in the treatment of gastrointestinal disorders, cardiovascular disorders and CNS disorders.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

PIPERIDINE DERIVATIVES AS 5-HT₄ RECEPTOR ANTAGONISTS

This invention relates to novel compounds having pharmacological activity, to
5 a process for their preparation and to their use as pharmaceuticals.

European Journal of Pharmacology 146 (1988), 187-188, and Naunyn-Schmiedeberg's Arch. Pharmacol. (1989) 340:403-410, describe a non classical 5-hydroxytryptamine receptor, now designated the 5-HT₄ receptor, and that ICS 205-930, which is also a 5-HT₃ receptor antagonist, acts as an antagonist at this
10 receptor.

WO 91/16045 (SmithKline and French Laboratories Limited) describes the use of cardiac 5-HT₄ receptor antagonists in the treatment of atrial arrhythmias and stroke.

EP-A-501322 (Glaxo Group Limited), WO 93/02677, WO 93/03725,
15 WO 93/05038, WO 93/05040 and WO 93/18036 (SmithKline Beecham plc) describe compounds having 5-HT₄ receptor antagonist activity.


It has now been discovered that certain novel compounds also have 5-HT₄ receptor antagonist properties.

Accordingly, the present invention provides compounds of formula (I) and
20 pharmaceutically acceptable salts thereof, and the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof:




wherein

25 X is a monocyclic or polycyclic aromatic group, such as a group of formula (a), (b), (c), (d), (e), (f) or (g):




(a)


- 2 -




(b)




(c)




(d)



(e)



(f)



(g)

wherein

L is N or CR_S wherein R_S is hydrogen, C₁₋₆ alkoxy, halogen, C₁₋₄ alkyl or cyano;

Q is NR_{1^a}, CH₂, O or S;

5 W is CH or N;

- in which $X_1-(CH_2)_x-X_2$ forms a 5-7 membered ring wherein X_1 is O or S; X_2 is O, S, NR or NRCO wherein R is hydrogen or C_{1-6} alkyl; and
 x is 1, 2 or 3;
 one of X_3 and X_4 is N and the other is C; and
- 5 X_5 is N or CR wherein R is hydrogen, C_{1-6} alkoxy, halo, C_{1-6} alkyl or cyano;
 R_1^a is hydrogen, C_{1-10} alkyl, C_{2-6} alkenyl, aralkyl, C_{2-6} alkanoyl or C_{2-6} alkanoyl C_{1-3} alkyl;
 R_3^a is hydrogen, halo, C_{1-6} alkyl, amino, nitro or C_{1-6} alkoxy;
 R_4^a is hydrogen, halo, C_{1-6} alkyl or C_{1-6} alkoxy;
- 10 R_1^b is C_{1-6} alkoxy; and
 R_2^b is hydrogen, chloro or fluoro;
 R_3^b is hydrogen, C_{1-6} alkyl, amino optionally substituted by a C_{1-6} alkyl group, halo, hydroxy or C_{1-6} alkoxy;
 R_4^b is hydrogen, halo, C_{1-6} alkyl, C_{1-6} alkoxy, nitro, amino or C_{1-6} alkylthio; and
- 15 R_5^b is hydrogen, halo, C_{1-6} alkyl, C_{1-6} alkoxy or amino;
 R_c is hydrogen, C_{1-6} alkoxy, halo or C_{1-6} alkyl;
 R_1^d is hydrogen, amino, halo, C_{1-6} alkyl, hydroxy or C_{1-6} alkoxy;
 R_2^d is hydrogen, halo, C_{1-6} alkyl, C_{1-6} alkoxy, nitro, amino or C_{1-6} alkylthio;
 R_3^d is hydrogen, halo, C_{1-6} alkyl, C_{1-6} alkoxy or amino;
- 20 R_4^d and R_5^d are independently hydrogen or C_{1-6} alkyl;
 R_1^e is hydrogen, halogen, CF_3 , C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} alkylthio, C_{1-6} alkylsulphonyl, C_{1-6} alkylsulphinyl, C_{1-7} acyl, cyano, C_{1-6} alkoxycarbonyl, C_{1-7} acylamino, hydroxy, nitro or amino, aminocarbonyl, or aminosulphonyl, optionally N-substituted by one or two groups selected from C_{1-6} alkyl, C_{3-8} cycloalkyl, and C_{3-8} cycloalkyl C_{1-4} alkyl or disubstituted by C_4 or C_5 polymethylene; phenyl or phenyl C_{1-4} alkyl group optionally substituted in the phenyl ring by one or two of halogen, C_{1-6} alkoxy or C_{1-6} alkyl groups;
- 25 R_3^e is hydrogen, halo, C_{1-6} alkyl, amino, nitro or C_{1-6} alkyl;
 R_4^e is hydrogen, halo, C_{1-6} alkyl or C_{1-6} alkoxy;
- 30 X_6-X_7 is NR_z-CO or $CR_1^fR_2^f-CR_3^fR_4^f$ where
 R_z and R_1^f to R_4^f are independently hydrogen or C_{1-6} alkyl; and/or
 R_1^f/R_2^f and R_3^f/R_4^f together are a bond and/or $R_1^f/R_2^f/R_3^f/R_4^f$ are joined to form C_{3-6} polymethylene;
- R_a^f is hydrogen, halo, C_{1-6} alkyl, amino, nitro or C_{1-6} alkyl;
- 35 R_b^f is hydrogen, halo, C_{1-6} alkyl or C_{1-6} alkoxy;
 X^g is O, S, SO, SO_2 , CH_2 , CH, N or NR wherein R is hydrogen or C_{1-6} alkyl;
 A is a saturated or unsaturated polymethylene chain of 2 - 4 carbon atoms;
 R_1^g and R_2^g are hydrogen or C_{1-6} alkyl;

- 4 -


R_3^g is hydrogen, halo, C_{1-6} alkyl, amino, nitro or C_{1-6} alkoxy;

R_4^g is hydrogen, halo, C_{1-6} alkyl or C_{1-6} alkoxy;

Y is O or NH;

Z is of sub-formula:

5



wherein

$-(CH_2)_n^1$ is attached at carbon; and

10 n^1 is 0, 1, 2, 3 or 4;

q is 0, 1, 2 or 3;

R_a is straight or branched chain alkylene of chain length 1-6 carbon atoms terminally substituted by R_7 wherein R_7 is aryl, 3 to 8 membered cycloalkyl, 3 to 8 membered heterocyclyl, 5 or 6 membered monocyclic heteroaryl or 9 or 10 membered fused bicyclic heteroaryl linked through carbon, or R_7 is C_{2-7} alkoxy carbonyl or secondary or tertiary hydroxy substituted C_{1-6} alkyl; and

15

R_6 is hydrogen or C_{1-6} alkyl;

or a compound of formula (I) wherein the CO-Y linkage is replaced by a heterocyclic bioisostere;

20

in the manufacture of a medicament for use as a 5-HT₄ receptor antagonist.

Examples of alkyl or alkyl containing groups include C_1 , C_2 , C_3 , C_4 , C_5 , C_6 , C_7 , C_8 , C_9 , C_{10} , C_{11} or C_{12} branched, straight chained or cyclic alkyl, as appropriate. C_{1-4} alkyl groups include methyl, ethyl, *n*- and *iso*-propyl, *n*-, *iso*-, *sec*- and *tert*-butyl. Cyclic alkyl includes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl optionally substituted by one of more alkyl groups of up to 4 carbon atoms.


25

Aryl includes phenyl and naphthyl optionally substituted by one or more substituents selected from halo, C_{1-6} alkyl and C_{1-6} alkoxy.

30

Halo includes fluoro, chloro, bromo and iodo, preferably chloro.

A suitable bioisostere for the amide or ester linkage containing Y in formula (I), is of formula:



wherein

the dotted circle represents one or two double bonds in any position in the 5-membered ring; H, J and I independently represent oxygen, sulphur, nitrogen or carbon, provided that at least one of H, J and I is other than carbon; U represents nitrogen or carbon.

- 5 Suitable examples of bioisosteres are as described for X, Y and Z in EP-A-328200 (Merck Sharp & Dohme Ltd.), such as an oxadiazole moiety.

L in formula (a) is favourably C-H, C-CH₃, C-Cl or C-OCH₃.

Q in formula (a) is favourably NR₁^a.

- 10 R₁^a is preferably hydrogen or a methyl or ethyl group.

R₁^b is preferably methoxy.

R₃^b is preferably amino.

R₄^b is preferably halo.

- 15 R₅^b is preferably hydrogen.

A substituent when halo is selected from fluoro, chloro, bromo and iodo.

R₄^a when halo is preferably iodo.

- 20 Suitable examples of the X₁-(CH₂)_x-X₂ moiety include O-(CH₂)₂-O, O-(CH₂)₃-O, O-CH₂-O, O-(CH₂)₂-NR, O-(CH₂)₂-S or O-CH₂-CONR, wherein any of the methylene linkages are optionally mono- or di- substituted by C₁₋₆ alkyl groups, such as methyl. Preferably X₁-(CH₂)₂-X₂ is O-(CH₂)₂-O.

R₁^d is preferably hydrogen or amino.

R₂^d is preferably hydrogen or halo.

- 25 R₃^d is preferably hydrogen or halo.

R₄^d and R₅^d are often hydrogen. When R₄^d/R₅^d is C₁₋₆ alkyl, it is often methyl. In particular R₄^d and R₅^d are methyl such that the disubstituent containing X₁ and X₂ is O-C(CH₃)₂-O.

- 30 R₁^e is preferably CF₃ or an ethyl group.

X₅ is preferably N, C-H or C-OCH₃;

R₃^e is preferably hydrogen.

R₄^e is preferably hydrogen or halo, such as iodo.

- 35 Suitable examples of X₆-X₇ when CR₁^fR₂^f-CR₃^fR₄^f include CH₂-CH₂ and CH=CH. X₆-X₇ is preferably NR₂-CO, however, such as NH-CO or N^tEt-CO.

R_a^f is preferably hydrogen.

R_b^f is preferably hydrogen or halo, such as iodo.

Values for A include $-\text{CH}_2-(\text{CH}_2)_r-\text{CH}_2-$ wherein r is 0, 1 or 2; $-\text{CH}_2-\text{CH}=\text{CH}-$; $-\text{C}(\text{CH}_3)=\text{CH}-$ or when X^g is CH or N, A may be $-(\text{CH}_2)_2-\text{CH}=-$ or $-\text{CH}=\text{CH}-\text{CH}=-$. Other examples of A are as described in the examples hereinafter.

- 5 R_1^g and R_2^g are often hydrogen or R_1^g and R_2^g are gem-dimethyl.
 r is often 1.
 R_3^g is preferably hydrogen.
 R_4^g is preferably hydrogen or halo, such as fluoro.

- 10 Other suitable values of X are as described in PCT/GB93/020208, PCT/EP93/02808, PCT/EP93/02775, PCT/EP93/02809, PCT/GB93/02130 (all in the name of SmithKline Beecham plc).

 Y is preferably O or NH.

- 15 n^1 is preferably 1 and the azacycle is preferably attached at a 4-position carbon atom, when q is 2.

 Values of Z of interest include 4-piperidinylmethyl and 4-pyrrolidinylmethyl, N-substituted by R_a .

- 20 Values for R_7 when monocyclic heteroaryl include pyridyl, pyrimidyl, pyrazinyl, pyrrol, imidazolyl, thienyl, furanyl, oxazole or thiazole (all possible isomers). Bicyclic heteroaryl R_7 include benzofuranyl, benzothiophenyl, indolyl and indazolyl, quinolyl and isoquinolyl (all possible isomers).

- Values for R_7 when 3 to 8 membered heterocyclyl, include cyclic
25 polymethylene interrupted by one or two of N, O or S, linked through C or N, for example N-linked piperidinyl or pyrrolidinyl.

- The pharmaceutically acceptable salts of the compounds of the formula (I) include acid addition salts with conventional acids such as hydrochloric, hydrobromic, boric, phosphoric, sulphuric acids and pharmaceutically acceptable
30 organic acids such as acetic, tartaric, maleic, citric, succinic, benzoic, ascorbic, methanesulphonic, α -keto glutaric, α -glycerophosphoric, and glucose-1-phosphoric acids.

- Examples of pharmaceutically acceptable salts include quaternary derivatives of the compounds of formula (I) such as the compounds quaternised by compounds
35 R_x-T wherein R_x is C_{1-6} alkyl, phenyl- C_{1-6} alkyl or C_{5-7} cycloalkyl, and T is a radical corresponding to an anion of an acid. Suitable examples of R_x include methyl, ethyl and *n*- and *iso*-propyl; and benzyl and phenethyl. Suitable examples of T include halide such as chloride, bromide and iodide.

Examples of pharmaceutically acceptable salts also include internal salts such as N-oxides.

The compounds of the formula (I), their pharmaceutically acceptable salts, (including quaternary derivatives and N-oxides) may also form pharmaceutically acceptable solvates, such as hydrates, which are included wherever a compound of formula (I) or a salt thereof is herein referred to.

The compounds of formula (I) wherein CO-Y is an ester or amide linkage are prepared by conventional coupling of the Z moiety with the appropriate acid. Suitable methods are as described in GB 2125398A (Sandoz Limited), GB 1593146A, EP-A-36269, EP-A-289170 and WO 92/05174 (Beecham Group p.l.c.). When CO-Y is replaced by a heterocyclic bioisostere, suitable methods are described in EP-A-328200 (Merck Sharp & Dohme Limited). Reference is also made to EP-A-501322 (Glaxo Group Limited).

The invention also comprises a process for preparing the compounds of formula (I) wherein X is of formula (d), which comprises reacting an appropriate benzoic acid derivative with an appropriate alcohol or amine. A process comprises reacting a benzoic acid derivative wherein the aromatic substituents are as required in the end compound of formula (I), or substituents convertible thereto, with an alcohol or amine containing Z or a group convertible thereto, and thereafter if necessary, converting the benzoic acid substituents and/or Z, and optionally forming a pharmaceutically acceptable salt.

Suitable examples of conversions in the aromatic substituents include chlorination of hydrogen to chloro, reduction of nitro to amino, dehydrohalogenation such as debromination, and/or elaboration of a 2,3-disubstituted benzoic acid with ethylene glycol to form the benzodioxan. Any elaboration of X is, however, usually carried out prior to ester or amide coupling.

Suitable examples of conversions in the Z containing moiety include conventional modifications of the N-substituent by substitution and/or deprotection or, in the case of a 2-, 3- or 4- substituted piperidinyll desired end compound, reduction of an appropriate pyridyl derivative.

Values for Z containing intermediates are as described in the aforementioned patent publications in the name of SmithKline Beecham plc.

The compounds of the present invention are 5-HT₄ receptor antagonists and it is thus believed may generally be used in the treatment or prophylaxis of gastrointestinal disorders, cardiovascular disorders and CNS disorders.

They are of potential interest in the treatment of irritable bowel syndrome (IBS), in particular the diarrhoea aspects of IBS, i.e., these compounds block the ability of 5-HT to stimulate gut motility via activation of enteric neurones. In animal

models of IBS, this can be conveniently measured as a reduction of the rate of defaecation. They are also of potential use in the treatment of urinary incontinence which is often associated with IBS.

They may also be of potential use in other gastrointestinal disorders, such as those associated with upper gut motility, and as antiemetics. In particular, they are of potential use in the treatment of the nausea and gastric symptoms of gastro-oesophageal reflux disease and dyspepsia. Antiemetic activity is determined in known animal models of cytotoxic-agent/radiation induced emesis.

Specific cardiac 5-HT₄ receptor antagonists which prevent atrial fibrillation and other atrial arrhythmias associated with 5-HT, would also be expected to reduce occurrence of stroke (see A.J. Kaumann 1990, Naumyn-Schmiedeberg's Arch. Pharmacol. 342, 619-622, for appropriate animal test method).

Anxiolytic activity is likely to be effected via the hippocampus (Dumuis *et al* 1988, Mol Pharmacol., 34, 880-887). Activity can be demonstrated in standard animal models, the social interaction test and the X-maze test.

Migraine sufferers often undergo situations of anxiety and emotional stress that precede the appearance of headache (Sachs, 1985, Migraine, Pan Books, London). It has also been observed that during and within 48 hours of a migraine attack, cyclic AMP levels are considerably increased in the cerebrospinal fluid (Welch *et al.*, 1976, Headache 16, 160-167). It is believed that a migraine, including the prodromal phase and the associated increased levels of cyclic AMP are related to stimulation of 5-HT₄ receptors, and hence that administration of a 5-HT₄ antagonist is of potential benefit in relieving a migraine attack.

Other CNS disorders of interest include schizophrenia, Parkinson's disease and Huntingdon's chorea.

The invention also provides a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

Such compositions are prepared by admixture and are usually adapted for enteral such as oral, nasal or rectal, or parenteral administration, and as such may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, nasal sprays, suppositories, injectable and infusible solutions or suspensions. Orally administrable compositions are preferred, since they are more convenient for general use.

Tablets and capsules for oral administration are usually presented in a unit dose, and contain conventional excipients such as binding agents, fillers, diluents, tableting agents, lubricants, disintegrants, colourants, flavourings, and wetting agents. The tablets may be coated according to well known methods in the art, for

example with an enteric coating.

Suitable fillers for use include cellulose, mannitol, lactose and other similar agents. Suitable disintegrants include starch, polyvinylpolypyrrolidone and starch derivatives such as sodium starch glycollate. Suitable lubricants include, for
5 example, magnesium stearate.

Suitable pharmaceutically acceptable wetting agents include sodium lauryl sulphate. Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid
10 preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminium stearate gel or hydrogenated edible fats, emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example, almond oil, fractionated
15 coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl *p*-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents.

Oral liquid preparations are usually in the form of aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs or are presented as a dry product
20 for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, emulsifying agents, non-aqueous vehicles (which may include edible oils), preservatives, and flavouring or colouring agents.

The oral compositions may be prepared by conventional methods of blending, filling or tableting. Repeated blending operations may be used to distribute the
25 active agent throughout those compositions employing large quantities of fillers. Such operations are, of course, conventional in the art.

For parenteral administration, fluid unit dose forms are prepared containing a compound of the present invention and a sterile vehicle. The compound, depending
30 on the vehicle and the concentration, can be either suspended or dissolved. Parenteral solutions are normally prepared by dissolving the compound in a vehicle and filter sterilising before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are also dissolved in the vehicle. To enhance the stability, the composition can be frozen after
35 filling into the vial and the water removed under vacuum.

Parenteral suspensions are prepared in substantially the same manner except that the compound is suspended in the vehicle instead of being dissolved and sterilised by exposure of ethylene oxide before suspending in the sterile vehicle.

- 10 -

Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound of the invention.

The invention further provides a method of treatment or prophylaxis of irritable bowel syndrome, dyspepsia, atrial arrhythmias and stroke, anxiety and/or migraine in mammals, such as humans, which comprises the administration of an effective amount of a compound of the formula (I) or a pharmaceutically acceptable salt thereof.

An amount effective to treat the disorders hereinbefore described depends on the relative efficacies of the compounds of the invention, the nature and severity of the disorder being treated and the weight of the mammal. However, a unit dose for a 70kg adult will normally contain 0.05 to 1000mg for example 0.5 to 500mg, of the compound of the invention. Unit doses may be administered once or more than once a day, for example, 2, 3 or 4 times a day, more usually 1 to 3 times a day, that is in the range of approximately 0.0001 to 50mg/kg/day, more usually 0.0002 to 25 mg/kg/day.

No adverse toxicological effects are indicated within the aforementioned dosage ranges.

The invention also provides a compound of formula (I) or a pharmaceutically acceptable salt thereof for use as an active therapeutic substance, in particular for use in the treatment of irritable bowel syndrome, gastro-oesophageal reflux disease, dyspepsia, atrial arrhythmias and stroke, anxiety and/or migraine.

The following Examples illustrates the preparation of compounds of formula (I), and the following Descriptions relate to the preparation of intermediates. The compounds of formula (I-1) and intermediates are prepared in Examples and Descriptions 1-1, 2-1 etc, the compounds of formula (I-2) are prepared in Examples and Descriptions 1-2, 2-2 etc and similarly for the compounds of formulae (I-3) to (I-5).

It will be appreciated that any compound prepared wherein Y is O may be provided as the corresponding compound wherein Y is NH.

A preferred compound corresponds to any of the compounds prepared in the Examples, but wherein there is an amino substituent in the 4-position and a chloro substituent in the 5-position of the benzoic acid nucleus depicted in formulae (I-1) to (I-5) inclusive.

Example 1 [X = (d), $X_1-(CH_2)_x-X_2 = O-(CH_2)_2-O$, $R_1^d = NH_2$, $R_2^d = Cl$, $R_3^d = H$, R_4^d , $R_5^d = H$; Y = O, Z = 4-piperidinylmethyl, $R_a = 3$ -pyridylmethyl]
5-(1-(3-Pyridylmethyl)-4-piperidinyl)methyl-8-amino-7-chloro-1,4-benzodioxancarboxylate (E1)

- 5 A stirred solution of 8-amino-7-chloro-1,4-benzodioxan-5-(4-piperidinylmethyl) carboxylate (0.1g, 0.31mmol) in acetone (10ml) was treated with Et_3N (0.043ml, 0.31mmol) and 3-picolyl chloride (0.043g, 0.34mmol). The reaction mixture was heated under reflux for 48 hours, cooled and evaporated *in vacuo*. The oily residue was purified by silica gel chromatography using $CHCl_3$ increasing to 2%
 10 MeOH, 98% $CHCl_3$ as eluant to yield the title compound as a pale yellow gum (0.05g) which was converted to the oxalate salt, mp 219 - 221°C.
 1H NMR 250MHz ($CDCl_3$) (free base)
 δ : 8.60-8.40(m, 2H), 7.65(d, 1H), 7.50(s, 1H), 7.30-7.20(m, 1H), 4.45(s, 2H), 4.40-4-
 25(m, 4H), 4.10(d, 2H), 3.50(s, 2H), 2.90(d, 2H), 2.00(t, 2H), 1.75(d, 2H),
 15 1.50-1.25(m, 3H)

Example 2 [X = (g), $X^g = O$, A = $(CH_2)_3$, (R_1^g , R_2^g , R_3^g , $R_4^g = H$; Y = NH, Z = 4-piperidinylmethyl, $R_a = 4$ -pyridylmethyl]
N-[(1-(4-Pyridylmethyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-
 20 **[1,3]oxazino[3,2-a]indole-10-carboxamide (E2)**

The title compound was prepared as an off-white solid by treating a solution of N-(4-piperidinylmethyl) 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide (D2) and triethylamine in acetonitrile and N,N-dimethylformamide with 4-picolyl chloride using the procedure described in Example 1.

- 25 1H NMR ($CDCl_3$)
 δ : 8.55(d, 2H), 8.35(d, 1H), 7.10-7.40(m, 5H), 6.57(t, 1H), 4.55(t, 2H), 4.14(t, 2H), 3.51(s, 2H), 3.37(t, 2H), 2.80-2.98(bd, 2H), 2.30-2.50(m, 2H), 1.93-2.14(m, 2H), 1.50-1.93(m, 3H), 1.31-1.50(m, 2H).

30 **Example 3** [X = (g), $X^g = O$, A = $(CH_2)_3$, (R_1^g , R_2^g , R_3^g , $R_4^g = H$; Y = NH, Z = 4-piperidinylmethyl, $R_a = 2$ -(1-piperidinyl)ethyl]
N-[(1-(2-(1-Piperidinyl)ethyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-
[1,3]oxazino[3,2-a]indole-10-carboxamide (E3)

- 35 The title compound was prepared as an off-white solid by treating a solution of N-(4-piperidinylmethyl) 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide (D2) and triethylamine in acetonitrile and N,N-dimethylformamide with 1-(2-chloroethyl)piperidine using the procedure described in Example 1; mp 139-141°C.

¹H NMR (CDCl₃)

δ: 8.35(d,1H), 7.07-7.48(m,3H), 6.55(t,1H), 4.56(t,2H), 4.13(t,2H), 3.35(t,2H), 2.90-3.07(bd,2H), 2.28-2.60(m,10H), 1.95-2.19(bt,2H), 1.23-1.90(m,11H).

- 5 **Example 4** [X = (g), X^g = O, A = (CH₂)₃, (R₁^g, R₂^g, R₃^g, R₄^g = H; Y = NH, Z = 4-piperidinylmethyl, R_a = benzofuran-2-ylmethyl]
N-[(1-(Benzofuran-2-ylmethyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide (E4)

- 10 The title compound was prepared as an off-white solid by treating a solution of N-(4-piperidinylmethyl) 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide (D2) and triethylamine in acetonitrile and N,N-dimethylformamide with 2-(chloromethyl)benzofuran (Blicke et al, J. Amer. Chem. Soc., 1949, 71, 2856) using the procedure described in Example 1.

¹H NMR (CDCl₃)

- 15 δ: 8.32(d,1H), 7.40-7.62(m,2H), 7.02-7.37(m,5H), 6.61(s,1H), 6.55(t,1H), 4.49(t,2H), 4.06(t,2H), 3.73(s,2H), 3.35(t,2H), 2.95-3.15(bd,2H), 2.25-2.46(m,2H), 2.02-2.25(bt,2H), 1.33-1.90(m,5H).

- 20 **Example 5** [X = (g), X^g = O, A = (CH₂)₃, (R₁^g, R₂^g, R₃^g, R₄^g = H; Y = NH, Z = 4-piperidinylmethyl, R_a = quinolin-2-ylmethyl)]
N-[(1-(Quinolin-2-ylmethyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide (E5)

- 25 The title compound was prepared by treating a solution of N-(4-piperidinylmethyl) 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide (D2) and triethylamine in acetonitrile and N,N-dimethylformamide with 2-(chloromethyl)quinoline using the procedure described in Example 1.

¹H NMR (CDCl₃)

- 30 δ: 8.32(d,1H), 8.16(d,1H), 8.08(d,1H), 7.65-7.90(m,3H), 7.54(t,1H), 7.07-7.35(m,3H), 6.58(t,1H), 4.52(t,2H), 4.09(t,2H), 3.96(s,2H), 3.35(t,2H), 3.00-3.15(m,2H), 2.25-2.50(m,4H), 1.35-1.95(m,5H).

Example 6 [X = (g), X^g = O, A = (CH₂)₃, (R₁^g, R₂^g, R₃^g, R₄^g = H; Y = NH, Z = 4-piperidinylmethyl, R_a = 5-phenylpentyl]

N-[(1-(5-Phenylpentyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide (E6)

5 The title compound was prepared by treating a solution of N-(4-piperidinylmethyl) 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide (D2) and triethylamine in acetonitrile and N,N-dimethylformamide with 1-chloro-5-phenylpentane using the procedure described in Example 1.

¹H NMR (CDCl₃)

10 δ: 8.28(d,1H), 7.08-7.30(m,8H), 6.61(t,1H), 4.54(t,2H), 4.10(t,2H), 3.34(t,2H), 3.15-3.27(bd,2H), 2.52-2.66(m,4H), 2.26-2.43(m,4H), 1.55-1.97(m,9H), 1.26-1.42(m,2H).

Example 7 [X = (g), X^g = O, A = (CH₂)₄, (R₁^g, R₂^g, R₃^g, R₄^g = H; Y = NH, Z = 4-piperidinylmethyl, R_a = 2-thienylmethyl]

15 **N-[(1-(2-Thienylmethyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide (E7)**

The title compound was prepared by treating a solution of N-(4-piperidinylmethyl) 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide (D2) and triethylamine in acetonitrile and N,N-dimethylformamide with 2-(chloromethyl)thiophene using the procedure described in Example 1.

¹H NMR(CDCl₃)

20 δ: 8.32(d,1H), 6.85-7.35(m,6H), 6.55(bt,1H), 4.52(t,2H), 4.10(t,2H), 3.74(s,2H), 3.32(t,2H), 2.90-3.05(m,2H), 2.25-2.50(m,4H), 1.30-1.90(m,5H).

25 **Example 8** [X = (g), X^g = O, A = (CH₂)₄, (R₁^g, R₂^g, R₃^g, R₄^g = H; Y = NH, Z = 4-piperidinylmethyl, R_a = 2-(cyclohexyl)ethyl]

N-[(1-(2-(Cyclohexyl)ethyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide (E8)

30 The title compound was prepared by treating a solution of N-(4-piperidinylmethyl) 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide (D2) and triethylamine in acetonitrile and N,N-dimethylformamide with 2-cyclohexylethyl bromide using the procedure described in Example 1.

¹H NMR(CDCl₃)

35 δ: 8.28(d,1H), 7.10-7.35(m,3H), 6.64(t,1H), 4.57(t,2H), 4.13(t,2H), 3.22-3.44(m,4H), 2.65-2.80(m,2H), 2.30-2.55(m,4H), 1.55-2.05(m,12H), 1.10-1.35(m,4H), 0.85-1.05(m,2H).

Example 9 [X = (g), X^g = O, A = (CH₂)₄, (R₁^g, R₂^g, R₃^g, R₄^g = H; Y = NH, Z = 4-piperidinylmethyl, R_a = 1-naphthylmethyl]

N-[(1-(1-Naphthylmethyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide (E9)

5 The title compound was prepared by treating a solution of N-(4-piperidinylmethyl) 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide (D2) and triethylamine in acetonitrile and N,N-dimethylformamide with 1-bromomethylnaphthylene using the procedure described in Example 1.

¹H NMR (CDCl₃)

10 δ: 8.23-8.48(m,2H), 7.70-7.90(m,2H), 7.33-7.57(m,4H), 7.00-7.30(m,3H), 6.52(t,1H), 4.46(t,2H), 4.01(t,2H), 3.90(s,2H), 3.31(t,2H), 2.90-3.07(bd,2H), 2.22-2.40(m,2H), 2.07(bt,2H), 1.55-1.85(m,3H), 1.20-1.50(m,2H).

Example 10 [X = (d), X₁-(CH₂)_x-X₂ = O-(CH₂)₂-O, R₁^d = NH₂, R₂^d = Cl, R₃^d = H, R₄^d, R₅^d = H; Y = O, Z = 4-piperidinylmethyl, R_a = 2-carboethoxyethyl]

15 **[1-(2-Carboethoxyethyl)-4-piperidinyl]methyl-8-amino-7-chloro-1,4-benzodioxan-5-carboxylate (E10)**

4-Piperidinylmethyl-8-amino-7-chloro-1,4-benzodioxan-5-carboxylate (0.100g; 0.31mmol) was dissolved in acetone (10ml) and treated with ethyl acrylate (0.037ml, 0.34mmol). The solution was heated at reflux (18 hours), cooled and evaporated *in vacuo* to a dark brown gum. The gum was purified by flash silica gel chromatography with CHCl₃→2% MeOH/CHCl₃ as eluant to yield the title compound as a colourless oil (0.053g; 41%) which was covered to the oxalate salt, mp=176-178°C.

25 ¹H NMR (270MHz, CD₃OD) (oxalate salt)

δ:7.40 (s, 1H), 4.27 (s, 2H), 4.20-4.10 (m, 2H), 3.60-3.50 (d, 2H), 3.37 (t, 2H), 3.27-3.23 (m, 6H), 3.10-2.90 (t, 2H), 2.85 (t, 2H), 2.10-1.90 (d, 2H), 1.70-1.55 (m, 1H), 1.30-1.20 (m, 3H)

30 **Example 11** [X = (d), X₁-(CH₂)_x-X₂ = O-(CH₂)₂-O, R₁^d = NH₂, R₂^d = Cl, R₃^d = H, R₄^d, R₅^d = H; Y = O, Z = 4-piperidinylmethyl, R_a = 3-hydroxybutyl]

[1-(3-Hydroxybutyl)-4-piperidinyl]methyl-8-amino-7-chloro-1,4-benzodioxan-5-carboxylate (E11)

a) 4-Piperidinylmethyl 8-amino-7-chloro-1,4-benzodioxan-5-carboxylate (0.100g; 0.31mmol) was dissolved in acetone (10ml) and treated with triethylamine (0.043ml; 0.31mmol) and methyl vinyl ketone (0.026ml; 0.34mmol). The solution was heated at reflux (18 hours), cooled and evaporated *in vacuo* to a yellow gum. The gum was purified by flash silica-gel chromatography with CHCl₃ as eluant to

yield [1-(3-oxobutyl)-4-piperidinyl]methyl 8-amino-7-chloro-1,4-benzodioxan-5-carboxylate as a colourless gum (0.050g; 41%) which was converted to the oxalate salt.

m.p. 160°C (Dec)

5 ¹H NMR (250 MHz, CDCl₃) (Free base)

δ: 7.47 (s, 1H), 4.47 (s, 2H), 4.40-4.30 (m, 4H), 4.10 (d, 2H), 2.90 (d, 2H), 2.65 (s, 4H), 2.17 (s, 3H), 2.00 (t, 2H), 1.85-1.70 (m, 2H), 1.47-1.25 (m, 3H)

b) [1-(3-oxobutyl)-4-piperidinyl]methyl-8-amino-7-chloro-1,4-benzodioxan-5-carboxylate (0.135g, 0.340mmol) (was dissolved in EtOH (8ml) and treated with
10 NaBH₄ (0.013g, 0.340mmol) with stirring. After 1h, the reaction mixture was evaporated under reduced pressure and the residue partitioned between CH₂Cl₂ and water. The aqueous layer was then extracted with CH₂Cl₂, and the combined organic layers were dried (Na₂SO₄) and evaporated under reduced pressure to give a colourless oil, which was purified by silica-gel chromatography (CH₂Cl₂/10% MeOH as eluant) to give the title compound as a colourless oil (0.048g, 35%), which was
15 converted to its oxalate salt m.p. 220-220°C.

¹H NMR (200MHz, CDCl₃) (free base) δ 7.50 (s, 1H), 4.50 (s, 2H), 4.35 (s, 4H), 4.10 (d, 2H), 3.96 (m, 1H), 3.25 (d, 1H), 3.00 (d, 1H), 2.65 (m, 2H), 2.18 (t, 1H), 2.00-1.35 (m, 9H), 1.20 (d, 3H).

20

Descriptions

Description 1

(1-Benzyl-4-piperidinyl)methylamine(D1)

25 A stirred solution of isonipecotamide (30.1g, 0.23 mole) and benzyl bromide (27.9 ml, 0.23 mole) in ethanol (250 ml) was treated with anhydrous potassium carbonate (64.9g, 0.47 mole) and heated under reflux for 3h. The mixture was allowed to cool, then filtered and the filtrate concentrated under vacuum. The residual oil was dissolved in chloroform (200 ml) and washed with water (1 x 150
30 ml), then dried (Na₂SO₄) and concentrated under vacuum to leave a yellow solid (41.0g). This solid was mixed thoroughly with phosphorus pentoxide (38.3g, 0.27 mole) and the mixture heated at 180°C under nitrogen for 2.5h with gentle stirring. The reaction mixture was allowed to cool, then treated with water (300 ml). When the solid mass had dissolved, the solution was basified by addition of solid K₂CO₃
35 and extracted with ethyl acetate (2x250 ml). The combined extracts were dried (Na₂SO₄) and concentrated *in vacuo* to leave a brown oil (35.3g). This was dissolved in dry ether (250 ml) and added dropwise over 30 minutes to a stirred suspension of lithium aluminium hydride (10.1g, 0.26 mole) in ether (150ml) at 0°C

under nitrogen. When addition was complete, the mixture was allowed to warm up to room temperature and was stirred for 1.5h. It was re-cooled to 0°C and treated cautiously with water (10ml), 10% NaOH solution (15 ml) and water again (25ml). The mixture was filtered through kieselguhr and the filtrate concentrated *in vacuo* to leave a brown oil, which was distilled under vacuum to afford the title compound as a colourless oil after distillation (27.8g, 67%) bp 106°C at 0.25 mmHg.

¹H NMR (CDCl₃)

δ: 7.20-7.37(m,5H), 3.48(s,2H), 2.85-2.95(m,2H), 2.55(d,2H), 1.87-2.00(m,2H), 1.60-1.75(m,2H), 1.10-1.40(m,5H).

Description 2

a) N-[(1-Benzyl-4-piperidinyl)methyl] indole-3-carboxamide

To a stirred solution of indole-3-carboxylic acid (15g, 0.093mole) in dichloromethane (250 ml) under nitrogen was added oxalyl chloride (8.7 ml, 0.10 mole) and dry dimethylformamide (6 drops). After 2 hours, the solvent was evaporated under reduced pressure. The residual acid chloride (0.093 mole) was dissolved in dichloromethane (100 ml) and added dropwise to a stirred solution of N-(1-benzyl-4-piperidinyl)methylamine (D1, 16.4g, 0.093 mole) and triethylamine (15.5 ml, 0.11 mole) in dichloromethane (150 ml) at 5°C. After stirring at ambient temperature overnight, the reaction mixture was washed with 10% Na₂CO₃ and the organic phase was dried (Na₂SO₄). The solvent was evaporated under reduced pressure and the residual solid recrystallised from ethyl acetate to afford the title compound as a white solid (17.5g, 60%).

¹H NMR (CDCl₃)

δ: 9.90(s,1H), 7.85-7.95(m,1H), 7.64(d,1H), 7.15-7.43(m,8H), 6.17(t,1H), 3.48(s,2H), 3.37(t,2H), 2.83-2.98(m,2H), 1.87-2.08(m,2H), 1.54-1.82(m,3H), 1.23-1.50(m,2H).

b) N-[(1-Benzyl-4-piperidinyl)methyl] 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide

A stirred suspension of N-[(1-benzyl-4-piperidinyl)methyl] indole-3-carboxamide (17.5g, 0.050 mole) in chloroform (250 ml) was treated with 3-bromo-1-propanol (10.1 ml, 0.11 mole) and N-chlorosuccinimide (8.7g, 0.065 mole) at room temperature and a clear solution was obtained in 15 minutes. After 1h the reaction mixture darkened in colour from pale yellow to orange and temperature rose to 38°C. After a further 1 h the reaction mixture was treated with 10% NaHCO₃ solution and the chloroform layer separated, dried (Na₂SO₄) and concentrated *in vacuo* to leave a yellow oil, which was chromatographed on silica gel eluting with 3% methanol/chloroform. The 2-(3-bromopropoxy)indole intermediate was dissolved in

acetone (400 ml), treated with anhydrous potassium carbonate (11g, 0.08 mole) and stirred at room temperature for 20h. The reaction mixture was concentrated *in vacuo* and the residue treated with water (200 ml) and extracted with chloroform (2 x 250 ml). The combined extracts were dried (Na₂SO₄), concentrated *in vacuo* and the residue chromatographed on silica gel eluting with 5% methanol/chloroform to afford the title compound as a pale yellow oil (3.1g, 15%). This was converted to its oxalate salt and crystallised from acetone as a white solid mp 169-170°C.

Free base:- ¹H NMR (CDCl₃)

δ: 8.32(d,1H), 7.05-7.38(m,8H), 6.53(t,1H), 4.50(t,2H), 4.08(t,2H), 3.48(s,2H), 3.31(t,2H), 2.83-2.97(m,2H), 2.27-2.41(m,2H), 1.54-2.06(m,5H), 1.25-1.45(m,2H).

c) N-(4-Piperidinylmethyl) 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide

A stirred suspension of N-[(1-benzyl-4-piperidinyl)methyl] 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide oxalate salt (2.25g, 0.0046 mole) in ethanol (100 ml) and glacial acetic acid (4 ml) was hydrogenated over 10% Pd-C (0.8g) at atmospheric pressure and 45°C for 18h. The mixture was filtered and the filtrate concentrated *in vacuo*. The majority of the product was in the solid which had been filtered off. This material was shaken with concentrated potassium carbonate solution (50 ml) and chloroform (50 ml) together with the residue from the filtrate. The mixture was filtered, the chloroform layer separated and dried (Na₂SO₄), then concentrated *in vacuo* to afford the title compound (D2) as a white solid (1.52g, 100%). This was recrystallised from chloroform/60-80 petrol mp 139-141°C.

¹H NMR (CDCl₃)

δ: 8.32(d,1H), 7.03-7.30(m,3H), 6.53(t,1H), 4.48(t,2H), 4.05(t,2H), 3.30(t,2H), 3.02-3.15(m,2H), 2.52-2.70(m,2H), 2.27-2.40(m,2H), 1.65-1.90(m,4H), 1.10-1.30(m,2H).

Description 3

4-Piperidinylmethyl 8-amino-7-chloro-1,4-benzodioxan-5-carboxylate hydrochloride

a) To a stirred solution of 8-amino-7-chloro-1,4-benzodioxan-5-carboxylic acid (prepared from the corresponding 7-H acid (prepared as in GB 1571278) by chlorination of the protected form) (1.10g) in acetonitrile was added bis-carbonyldiimidazole (0.77g). The reaction mixture was stirred at room temperature for 1.5 hours. The solvent was removed under reduced pressure to afford crude 8-amino-7-chloro-1,4-benzodioxan-5-imidazolide.

b) To a solution of N-*tert*-butoxycarbonyl-4-hydroxymethyl piperidine (0.25g) in dry THF (10ml) was added methyllithium (1.5M in diethylether; 0.78ml) at 0°C under a nitrogen atmosphere. Stirring was continued at ambient temperature for 10

- 18 -

min. 8-Amino-7-chloro-1,4-benzodioxan-5-imidazolid (0.33g) in THF (10ml) was added to the reaction mixture and stirring continued for 2 hours. The reaction mixture was cooled to 0°C and water was added. The solvent was removed under reduced pressure and the residue partitioned between chloroform and water. The organic phase was washed with water (3x), dried (Na₂SO₄) filtered and concentrated *in vacuo*. Flash chromatography on silica using chloroform and ethanol as eluant gave the title compound (0.26g).

¹H NMR 250MHz (CDCl₃)

δ: 7.47(s,1H), 4.49(s,2H), 4.36(s,4H), 4.08-4.22(m,4H), 2.64-2.80(m,2H), 1.84-2.01(m,1H), 1.70-1.83(m,2H), 1.46(s,9H), 1.18-1.38(m,2H)

c) HCl(g) was bubbled into a cooled solution of 8-amino-7-chloro-(N-*tert*-butoxycarbonyl-4-piperidylmethyl)-1,4-benzodioxan-5-carboxylate (0.26g) in dioxan (50ml) for 25 min. The solvent was concentrated *in vacuo* and the residue triturated with Et₂O to afford pure title compound (0.12g).

mp 249-251°C

¹H NMR 250MHz (DMSO)

δ: 8.99-9.10(m,1H), 8.59-8.78(m,1H), 7.29(s,1H), 5.73(s,2H), 4.25-4.34(s,4H), 4.03(d,2H), 3.20-3.42(m,2H), 2.75-2.97(m,2H), 1.76-2.06(m,3H), 1.48-1.57(m,2H)

5-HT₄ RECEPTOR ANTAGONIST ACTIVITY

1) Guinea pig colon

Male guinea-pigs, weighing 250-400g are used. Longitudinal muscle-myenteric plexus preparations, approximately 3cm long, are obtained from the distal colon region. These are suspended under a 0.5g load in isolated tissue baths containing Krebs solution bubbled with 5% CO₂ in O₂ and maintained at 37°C. In all experiments, the Krebs solution also contains methiothepin 10⁻⁷M and granisetron 10⁻⁶M to block effects at 5-HT₁, 5-HT₂ and 5-HT₃ receptors.

After construction of a simple concentration-response curve with 5-HT, using 30s contact times and a 15min dosing cycle, a concentration of 5-HT is selected so as to obtain a contraction of the muscle approximately 40-70% maximum (10⁻⁹M approx). The tissue is then alternately dosed every 15min with this concentration of 5-HT and then with an approximately equi-effective concentration of the nicotine receptor stimulant, dimethylphenylpiperazinium (DMPP). After obtaining consistent responses to both 5-HT and DMPP, increasing concentrations of a putative 5-HT₄ receptor antagonist are then added to the bathing solution. The effects of this compound are then determined as a percentage reduction of the contractions evoked by 5-HT or by DMPP. From this data, pIC₅₀ values are determined, being defined as the -log concentration of antagonist which reduces the contraction by 50%. A

compound which reduces the response to 5-HT but not to DMPP is believed to act as a 5-HT₄ receptor antagonist.

The compounds tested had a pIC₅₀ of >7, E1 had a pIC₅₀ of >9.

2) Rat oesophagus

- 5 Rat oesophageal tunica muscularis mucosae is set up according to Baxter *et al.* Naunyn-Schmiedeberg's Arch. Pharmacol., 343, 439-446 (1991). The inner smooth muscle tube of the muscularis mucosae is isolated and mounted for isometric tension recording in oxygenated (95% O₂/5% CO₂) Tyrodes solution at 37°C. All experiments are performed in pargyline pre-treated preparations (100µM for 15 min
10 followed by washout) and in the presence of cocaine (30µM). Relaxant responses to 5-HT are obtained after pre-contracting the oesophagus tissue with carbachol (3µM).

Claims

1. Compounds of formula (I) and pharmaceutically acceptable salts thereof, and the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof:



5




wherein


X is a monocyclic or polycyclic aromatic group, such as a group of formula (a), (b), (c), (d), (e), (f) or (g):

10






(e)



(f)



(g)

wherein

L is N or CR_s wherein R_s is hydrogen, C₁₋₆ alkoxy, halogen, C₁₋₄ alkyl or cyano;

Q is NR_{1^a}, CH₂, O or S;

5 W is CH or N;

in which X₁-(CH₂)_x-X₂ forms a 5-7 membered ring wherein X₁ is O or S; X₂ is O, S, NR or NRCO wherein R is hydrogen or C₁₋₆ alkyl; and
x is 1, 2 or 3;

one of X₃ and X₄ is N and the other is C; and

10 X₅ is N or CR wherein R is hydrogen, C₁₋₆ alkoxy, halo, C₁₋₆ alkyl or cyano;

R_{1^a} is hydrogen, C₁₋₁₀ alkyl, C₂₋₆ alkenyl, aralkyl, C₂₋₆ alkanoyl or C₂₋₆ alkanoyl
C₁₋₃ alkyl;

R_{3^a} is hydrogen, halo, C₁₋₆ alkyl, amino, nitro or C₁₋₆ alkoxy;

R_{4^a} is hydrogen, halo, C₁₋₆ alkyl or C₁₋₆ alkoxy;

15 R_{1^b} is C₁₋₆ alkoxy; and

R_{2^b} is hydrogen, chloro or fluoro;

R_{3^b} is hydrogen, C₁₋₆ alkyl, amino optionally substituted by a C₁₋₆ alkyl group,
halo, hydroxy or C₁₋₆ alkoxy;


R_{4^b} is hydrogen, halo, C₁₋₆ alkyl, C₁₋₆ alkoxy, nitro, amino or C₁₋₆ alkylthio; and

20 R_{5^b} is hydrogen, halo, C₁₋₆ alkyl, C₁₋₆ alkoxy or amino;

R_c is hydrogen, C₁₋₆ alkoxy, halo or C₁₋₆ alkyl;

R_{1^d} is hydrogen, amino, halo, C₁₋₆ alkyl, hydroxy or C₁₋₆ alkoxy;

- R_2^d is hydrogen, halo, C_{1-6} alkyl, C_{1-6} alkoxy, nitro, amino or C_{1-6} alkylthio;
 R_3^d is hydrogen, halo, C_{1-6} alkyl, C_{1-6} alkoxy or amino;
 R_4^d and R_5^d are independently hydrogen or C_{1-6} alkyl;
 R_1^e is hydrogen, halogen, CF_3 , C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} alkylthio, C_{1-6}
 5 alkylsulphonyl, C_{1-6} alkylsulphinyl, C_{1-7} acyl, cyano, C_{1-6} alkoxycarbonyl,
 C_{1-7} acylamino, hydroxy, nitro or amino, aminocarbonyl, or aminosulphonyl,
 optionally N-substituted by one or two groups selected from C_{1-6} alkyl, C_{3-8}
 cycloalkyl, and C_{3-8} cycloalkyl C_{1-4} alkyl or disubstituted by C_4 or C_5
 polymethylene; phenyl or phenyl C_{1-4} alkyl group optionally substituted in
 10 the phenyl ring by one or two of halogen, C_{1-6} alkoxy or C_{1-6} alkyl groups;
 R_3^e is hydrogen, halo, C_{1-6} alkyl, amino, nitro or C_{1-6} alkyl;
 R_4^e is hydrogen, halo, C_{1-6} alkyl or C_{1-6} alkoxy;
 X_6-X_7 is NR_2-CO or $CR_1^fR_2^f-CR_3^fR_4^f$ where
 R_2 and R_1^f to R_4^f are independently hydrogen or C_{1-6} alkyl; and/or
 15 R_1^f/R_2^f and R_3^f/R_4^f together are a bond and/or $R_1^f/R_2^f/R_3^f/R_4^f$ are joined to form
 C_{3-6} polymethylene;
 R_a^f is hydrogen, halo, C_{1-6} alkyl, amino, nitro or C_{1-6} alkyl;
 R_b^f is hydrogen, halo, C_{1-6} alkyl or C_{1-6} alkoxy;
 X_8 is O, S, SO, SO_2 , CH_2 , CH, N or NR wherein R is hydrogen or C_{1-6} alkyl;
 20 A is a saturated or unsaturated polymethylene chain of 2 - 4 carbon atoms;
 R_1^g and R_2^g are hydrogen or C_{1-6} alkyl;
 R_3^g is hydrogen, halo, C_{1-6} alkyl, amino, nitro or C_{1-6} alkoxy;
 R_4^g is hydrogen, halo, C_{1-6} alkyl or C_{1-6} alkoxy;
 Y is O or NH;
 25 Z is of sub-formula:



- wherein
 30 $-(CH_2)_n^1$ is attached at carbon; and
 n^1 is 0, 1, 2, 3 or 4;
 q is 0, 1, 2 or 3;
 R_a is straight or branched chain alkylene of chain length 1-6 carbon atoms terminally
 substituted by R_7 wherein and R_7 is aryl, 3 to 8 membered cycloalkyl, 3 to 8
 35 membered heterocyclyl, 5 or 6 membered monocyclic heteroaryl or 9 or 10
 membered fused bicyclic heteroaryl linked through carbon, or R_7 is

C₂₋₇ alkoxy carbonyl or secondary or tertiary hydroxy substituted C₁₋₆ alkyl;
and

R₆ is hydrogen or C₁₋₆ alkyl;

or a compound of formula (I) wherein the CO-Y linkage is replaced by a heterocyclic

5 bioisostere;

in the manufacture of a medicament for use as a 5-HT₄ receptor antagonist.

2. A compound according to claim 1 wherein:

10 L in formula (a) is C-H, C-CH₃, C-Cl or C-OCH₃;

Q in formula (a) is NR₁^a;

R₁^a is hydrogen or a methyl or ethyl group.

3. A compound according to claim 1 wherein:

15 R₁^b is methoxy;

R₃^b is amino;

R₄^b is halo;

R₅^b is hydrogen.

4. A compound according to claim 1 wherein:

20 X₁-(CH₂)_x-X₂ moiety is O-(CH₂)₂-O, O-(CH₂)₃-O, O-CH₂-O, O-(CH₂)₂-NR,
O-(CH₂)₂-S or O-CH₂-CONR, wherein any of the methylene linkages are optionally
mono- or di- substituted by C₁₋₆ alkyl groups;

R₁^d is hydrogen or amino;

25 R₂^d is hydrogen or halo;

R₃^d is hydrogen or halo.

5. A compound according to claim 1 wherein:

R₁^e is CF₃ or an ethyl group;

30 X₅ is N, C-H or C-OCH₃;

R₃^e is hydrogen;

R₄^e is hydrogen or halo, such as iodo.

6. A compound according to claim 1 wherein:

35 X₆-X₇ when CR₁^fR₂^f-CR₃^fR₄^f is CH₂-CH₂, CH=CH; NH-CO or NEt-CO;

R_a^f is hydrogen;

R_b^f is hydrogen or halo.

7. A compound according to claim 1 wherein:
A is $-\text{CH}_2-(\text{CH}_2)_r-\text{CH}_2-$ wherein r is 0, 1 or 2; $-\text{CH}_2-\text{CH}=\text{CH}-$; $-\text{C}(\text{CH}_3)=\text{CH}-$ or
when X⁸ is CH or N, A may be $-(\text{CH}_2)_2-\text{CH}=-$ or $-\text{CH}=\text{CH}-\text{CH}=-$;
R₁⁸ and R₂⁸ are hydrogen or R₁⁸ and R₂⁸ are gem-dimethyl;
5 r is 1;
R₃⁸ is hydrogen;
R₄⁸ is hydrogen or halo.
8. A compound according to any one of claims 1 to 7 wherein Y is O or NH.
- 10 9. A compound according to any one of claims 1 to 8 wherein n¹ is 1 and the
azacycle is attached at a 4-position carbon atom, when q is 2, and Z is
4-piperidinylmethyl and 4-pyrrolidinylmethyl, N-substituted by R_a as defined in
claim 1.
- 15 10. A compound according to any one of claims 1 to 9 wherein Z is as in any of
the Examples hereinbefore described.
- 20 11. 5-(1-(3-Pyridylmethyl)-4-piperidinyl)methyl-8-amino-7-chloro-1,4-
benzodioxancarboxylate.
12. N-[(1-(4-Pyridylmethyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-
[1,3]oxazino[3,2-a]indole-10-carboxamide.
- 25 13. N-[(1-(2-(1-Piperidinyl)ethyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-
[1,3]oxazino[3,2-a]indole-10-carboxamide.
14. N-[(1-(Benzofuran-2-ylmethyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-
[1,3]oxazino[3,2-a]indole-10-carboxamide.
- 30 15. N-[(1-(Quinolin-2-ylmethyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-
[1,3]oxazino[3,2-a]indole-10-carboxamide.
16. N-[(1-(5-Phenylpentyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-
35 [1,3]oxazino[3,2-a]indole-10-carboxamide.
17. N-[(1-(2-Thienylmethyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-
[1,3]oxazino[3,2-a]indole-10-carboxamide.

18. N-[(1-(2-(Cyclohexyl)ethyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide.
- 5 19. N-[(1-(1-Naphthylmethyl)-4-piperidinyl)methyl] 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide.
20. [1-(2-Carboethoxyethyl)-4-piperidinyl]methyl-8-amino-7-chloro-1,4-benzodioxan-5-carboxylate.
- 10 21. [1-(3-Hydroxybutyl)-4-piperidinyl]methyl-8-amino-7-chloro-1,4-benzodioxan-5-carboxylate.
22. A compound according to any one of claims 7 to 10 in the form of a
15 pharmaceutically acceptable salt.
23. A compound according to any one of claims 10 to 22 but wherein Y is NH.
24. A process for preparing the ester or amide compounds (where Y is O or NH)
20 according to claim 1, which comprises reacting an appropriate acid derivative with an appropriate alcohol or amine.
25. A pharmaceutical composition comprising a compound according to any one of claims 1 to 23, and a pharmaceutically acceptable carrier.
- 25 26. A compound according to claim 1 for use as an active therapeutic substance.
27. The use of a compound according to claim 1 in the manufacture of a
30 medicament for use as a 5-HT₄ receptor antagonist.
28. The use according to claim 27 for use as a 5-HT₄ receptor antagonist in the treatment or prophylaxis of gastrointestinal disorders, cardiovascular disorders and CNS disorders.
- 35

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/EP 93/03054

A. CLASSIFICATION OF SUBJECT MATTER

IPC 5 C07D498/04 C07D405/14 C07D405/12 A61K31/535 A61K31/445
 //C07D498/04,265:00,209:00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB,A,2 176 785 (ASTRA LAKEMEDEL) 7 January 1987 see claims 1,12,16 ---	1,25,28
X	EP,A,0 407 137 (YOSHITOMI) 9 January 1991 see page 1; claims 1,5 ---	1,25,28
P,X	WO,A,93 03725 (SMITH KLINE BEECHAM) 4 March 1993 cited in the application see page 49 see page 2 - page 3; claims 1,10 ---	1,25,28
P,X	WO,A,93 05038 (SMITHKLINE BEECHAM) 18 March 1993 cited in the application see claims 1,11,16,17 ---	1,25,28
	--- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

9 February 1994

Date of mailing of the international search report

18. 02. 94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

Voyiazoglou, D

INTERNATIONAL SEARCH REPORT

Inter. Appl. Application No

PCT/EP 93/03054

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO,A,93 18036 (SMITHKLINE BEECHAM) 16 November 1993 cited in the application see claims 1,12,15 ---	1,25,28
A	EP,A,0 501 322 (GLAXO GROUP) 2 September 1992 cited in the application see claims 1,15,16 -----	1,25,28

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 93/03054

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB-A-2176785	07-01-87	NONE	
EP-A-0407137	09-01-91	CA-A- 2020315 US-A- 5185333 JP-A- 3279372	04-01-91 09-02-93 10-12-91
WO-A-9303725	04-03-93	NONE	
WO-A-9305038	18-03-93	AU-A- 2491092 AU-A- 2541892 WO-A- 9305040 AU-A- 2435092 AU-B- 3457293 WO-A- 9316072 WO-A- 9324117 WO-A- 9400113	05-04-93 05-04-93 18-03-93 16-03-93 03-09-93 19-08-93 09-12-93 06-01-94
WO-A-9318036	16-09-93	WO-A- 9324117	09-12-93
EP-A-0501322	02-09-92	AU-B- 645402 AU-A- 1209492 WO-A- 9214727	13-01-94 15-09-92 03-09-92